

# MATH 142A: Introduction to Analysis

[math-old.ucsd.edu/~ynemish/teaching/142a](http://math-old.ucsd.edu/~ynemish/teaching/142a)

Today: Uniform continuity  
> Q&A: February 11

Next: Ross § 19

Week 6:

- Homework 5 (due Sunday, February 13)

## Inverse function

Def 18.9 Function  $f: X \rightarrow Y$  is called **one-to-one** (or **bijection**)


$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$$

if  $f(x)=y$  and  $\forall y \in Y \exists! x \in X$  s.t.  $f(x)=y$

Example  $\sin: [-\frac{\pi}{2}, \frac{\pi}{2}] \rightarrow [-1, 1]$  is one-to-one

$\sin: [0, \pi] \rightarrow [0, 1]$  is not one-to-one

$$\sin(0) = \sin(\pi) = 0$$



Def 18.10 Let  $f: X \rightarrow Y$  be a bijection,  $y = f(x)$ .

Then the function  $f^{-1}: Y \rightarrow X$  given by  $(f^{-1}(y) = x \Leftrightarrow f(x) = y)$

is called the **inverse of  $f$** . In particular  $f^{-1}(f(x)) = x$ ,  $f(f^{-1}(y)) = y$

Example •  $\sin: [-\frac{\pi}{2}, \frac{\pi}{2}] \rightarrow [-1, 1]$ ,  $\sin^{-1} = \arcsin: [-1, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$

•  $f: [0, +\infty) \rightarrow [0, +\infty)$ ,  $f(x) = x^m$ ,  $f^{-1}: [0, +\infty) \rightarrow [0, +\infty)$ ,  $f^{-1}(x) = x^{\frac{1}{m}} = \sqrt[m]{x}$

• If  $f$  is strictly increasing (decreasing) on  $X$ , then  $f: X \rightarrow f(X)$  is a bijection

## Continuity and the inverse function

Thm 18.4 Let  $f$  be a continuous strictly increasing function on some interval  $I$ . Then  $J := f(I)$  is an interval and

$f^{-1}: J \rightarrow I$  is

Proof ①  $f^{-1}$  is strictly increasing: Take  $y_1, y_2 \in J$ ,  $y_1 < y_2$

Denote  $x_1 = f^{-1}(y_1)$ ,  $x_2 = f^{-1}(y_2)$ . Then

If  $x_1 \geq x_2$ , then

②  $J$  is an interval: By Cor. 18.3  $J$  is either an or

a Since  $f$  is strictly increasing,  $J$  is an

③ ① + ② +

## One-to-one continuous functions

Thm 18.6 Let  $f$  be a one-to-one continuous function on an interval  $I$ . Then  $f$  is or

Proof. ① If  $a < b < c$  then either or

Otherwise, or


If  $f(b) > \max\{f(a), f(c)\}$ , choose

Then by Thm 18.2

Similarly when  $f(b) < \min\{f(a), f(c)\}$ .

② Take any  $a_0 < b_0$ . If  $f(a_0) < f(b_0)$ , then  $f$  is on  $I$ .

③ Similarly, if  $f(a_0) > f(b_0)$ , then  $f$  is decreasing.



## Uniform continuity

Def. (Continuity on a set) Function  $f$  is continuous on  $S \subset \mathbb{R}$

if  $\forall x \in S \quad \forall \varepsilon > 0 \quad \exists \delta > 0 \quad \forall y \in S \quad \text{s.t.} \quad |x - y| < \delta \quad (|f(x) - f(y)| < \varepsilon)$

Def. (Uniform continuity) Function  $f$  is uniformly continuous on  $S \subset \mathbb{R}$  if

Example Let  $f(x) = \frac{1}{x}$ .

1)  $\forall [a,b] \subset (0, +\infty) \quad f$  is unif. cont. on  $[a,b]$ .

Fix  $\varepsilon > 0$ . Then for  $x, y \in [a, b]$


. Take

Then

2)  $f$  is not unif. cont. on  $(0, 1]$ . Fix

Then

, but



## Cantor - Heine Theorem

Remark If  $f$  is uniformly continuous on  $S \subset \mathbb{R}$ , then  $f$  is continuous on  $S$ .

Thm 19.2 If  $f$  is continuous on a closed interval  $[a, b]$ , then

$f$  is

Proof. Suppose that  $f$  is cont. but not unif. cont. on  $[a, b]$ .

$\Rightarrow$

Take

and thus

, so

## Uniform continuity

Thm 19.4 If  $f$  is uniformly continuous on a set  $S$ , and  $(s_n)$  is a Cauchy sequence in  $S$ , then  $(f(s_n))$  is a Cauchy sequence

Proof. Fix  $\epsilon > 0$ .

①  $f$  is unif. cont. on  $S$

②  $(s_n)$  is a Cauchy sequence

## Example

Consider  $f(x) = \frac{1}{x}$  and  $t_n = \frac{1}{n}$ .  $(t_n)$  is a Cauchy sequence,  $\forall n t_n \in (0, 1]$ , but  $f(t_n) = n$  is not a Cauchy sequence.  
 $\Rightarrow f$  is not unif. cont. on  $(0, 1]$ .

## Examples

3)  $f(x) = x^2$  is continuous on  $\mathbb{R}$ , but is not unif. continuous on  $\mathbb{R}$ .

Take a sequence

Then

(i)

(ii)

4)  $f(x) =$  is continuous and bounded on  $\mathbb{R}$ , but not  
unif. continuous on  $\mathbb{R}$

Take

Then

(i)

(ii)