

MATH180C: Introduction to Stochastic Processes II

<https://mathweb.ucsd.edu/~ynemish/teaching/180c>

Today: Introduction. Birth processes

Next: PK 6.2-6.3

Week 1:

- join Piazza

Continuous Time Markov Chains

Def (Continuous-time Markov chain)

Let $(X_t)_{t \geq 0} = (X_t : 0 \leq t < \infty)$ be a continuous time process taking values in \mathbb{Z}_+ . $(X_t)_{t \geq 0}$ is called Markov chain if for any $n \in \mathbb{N}$, $0 \leq t_0 < t_1 < \dots < t_{n-1} < s, t > 0, i_0, i_1, \dots, i_{n-1}, i, j \in \mathbb{Z}_+$

$$\begin{aligned} P(X_{s+t} = j | X_{t_0} = i_0, X_{t_1} = i_1, \dots, X_{t_{n-1}} = i_{n-1}, X_s = i) \\ = P(X_{s+t} = j | X_s = i) \end{aligned}$$

Transition probability function

One way of describing a continuous time MC is by using the transition probability functions.

Def. Let $(X_t)_{t \geq 0}$ be a MC. We call

$$i, j \in \{0, 1, \dots\}, s \geq 0, t > 0$$

the transition probability function for $(X_t)_{t \geq 0}$.

If $P(X_{s+t} = j | X_s = i)$ does not depend on s , we say that $(X_t)_{t \geq 0}$ has stationary transition probabilities and we define

[compare with n-step transition probabilities]

Characterization of the Poisson process

Experiment: count events occurring along $[0, +\infty)$ { or 1-D space



Denote by $N((a, b])$ the number of events that occur on $(a, b]$.

Assumptions:

1. Number of events happening in disjoint intervals are independent.
2. For any $t \geq 0$ and $h > 0$, the distribution of $N((t, t+h])$ does not depend on t (only on h , the length of the interval)
3. There exists $\lambda > 0$ s.t. $P(N((t, t+h]) \geq 1) = \lambda h + o(h)$ as $h \rightarrow 0$ (rare events)
4. Simultaneous events are not possible: $P(N((t, t+h]) \geq 2) = o(h), h \rightarrow 0$

Transition probabilities of the Poisson process

Let $(X_t)_{t \geq 0}$ be the Poisson process.

Define the transition probability functions

$$P_{ij}(h) := P(X_{t+h} = j | X_t = i), \quad i, j \in \{0, 1, 2, \dots\}, \quad t \geq 0, \quad h > 0$$

What are the infinitesimal (small h) transition probability functions for $(X_t)_{t \geq 0}$? As $h \rightarrow 0$,

$$P_{ii}(h) = P(X_{t+h} = i | X_t = i)$$

=

$$P_{i,i+1}(h) = P(X_{t+h} = i+1 | X_t = i) =$$

$$\sum_{j \notin \{i, i+1\}} P_{ij}(h) =$$

Poisson process and transition probabilities

To sum up: $(X_t)_{t \geq 0}$ is a MC with (infinitesimal) transition probabilities satisfying

$$P_{ii}(h) =$$

$$P_{i,i+1}(h) =$$

$$\sum_{j \notin \{i, i+1\}} P_{i,j}(h) =$$

What if we allow $P_{ij}(h)$ depend on i ?

↳ birth and death processes

Pure birth processes

Def Let $(\lambda_k)_{k \geq 0}$ be a sequence of positive numbers.

We define a pure birth process as a Markov process

$(X_t)_{t \geq 0}$ whose stationary transition probabilities satisfy

$$1. P_{k,k+1}(h) =$$

$$2. P_{k,k}(h) =$$

$$3. P_{k,j}(h) =$$

$$4. X_0 = 0$$

Related model. Yule process : $\lambda_k = \beta k$ for some $\beta > 0$.

Describes the growth of a population

- birth rate is proportional to the size of the population

Birth processes and related differential equations

Now define $P_n(t) = P(X_t = n)$. For small $h > 0$

$$P_n(t+h) = P(X_{t+h} = n) =$$

=

=

=

=

$$P_n(t+h) - P_n(t) = -\lambda_n h P_n(t) + \lambda_{n-1} h P_{n-1}(t) + o(h)$$

$$P_n'(t) =$$

Birth processes and related differential equations

$P_n(t)$ satisfies the following system

of differential eqs.


with initial conditions

$$(*) \quad \left\{ \begin{array}{l} P_0'(t) = \\ P_1'(t) = \\ P_2'(t) = \\ \vdots \\ P_n'(t) = \\ \vdots \end{array} \right. \quad \left\{ \begin{array}{l} P_0(0) = \\ P_1(0) = \\ P_2(0) = \\ \vdots \\ P_n(0) = \\ \vdots \end{array} \right.$$

Solving this system gives the p.m.f. of X_t for any t

Description of the birth processes via sojourn times

$$(X_t)_{t \geq 0}$$



w_i - i-th "birth time" s_i - "time between (i-1)-th birth and i-th birth"

$$w_i = \sum_{\ell=0}^{i-1} s_\ell \quad \hookrightarrow \text{sojourn times}$$

Alternative way of characterizing $(X_t)_{t \geq 0}$:

-

-

Description of the birth processes via sojourn times

Theorem

Let $(\lambda_k)_{k \geq 0}$ be a sequence of positive numbers. Let $(X_t)_{t \geq 0}$ be a non-decreasing right-continuous process, $X_0 = 0$, taking values in $\{0, 1, 2, \dots\}$. Let $(S_i)_{i \geq 0}$ be the sojourn times associated with $(X_t)_{t \geq 0}$, and define $W_e = \sum_{i=0}^{e-1} S_i$.

Then conditions

(a)



(b)

are equivalent to

(c)

Explosion

$(X_t)_{t \geq 0}$



population becomes infinite in finite time

Thm. Let $(X_t)_{t \geq 0}$ be a pure birth process of rates $(\lambda_k)_{k \geq 0}$.

Then

Solving the system of differential equations (*)

$$(*) \begin{cases} P_0'(t) = -\lambda_0 P_0(t), & P_0(0) = 1 \\ P_n'(t) = -\lambda_n P_n(t) + \lambda_{n-1} P_{n-1}(t), & P_n(0) = 0 \quad \text{for } n \geq 1 \end{cases}$$

$P_0(t)$:

$$P_0'(t) =$$

$$\frac{P_0'(t)}{P_0(t)} =$$

$$g'(t) =$$

$$g(t) =$$

Solving the system of differential equations (*)

$P_n(t)$, $n \geq 1$

Consider the function $Q_n(t) =$

$$(Q_n(t))' =$$

$$Q_n(t) =$$

$$\mathcal{L}_1 P_n(t) =$$

← apply recursively

$$\begin{aligned} P_1(t) &= e^{-\lambda_1 t} \int_0^t \lambda_0 e^{\lambda_1 s} e^{-\lambda_0 s} ds = e^{-\lambda_1 t} \int_0^t \lambda_0 e^{(\lambda_1 - \lambda_0)s} ds \quad (\text{if } \lambda_1 \neq \lambda_0) \\ &= e^{-\lambda_1 t} \frac{\lambda_0}{\lambda_1 - \lambda_0} \left(e^{(\lambda_1 - \lambda_0)t} - 1 \right) = \frac{\lambda_0}{\lambda_1 - \lambda_0} e^{-\lambda_1 t} - \frac{\lambda_0}{\lambda_1 - \lambda_0} e^{-\lambda_1 t} \end{aligned}$$